

UATP-95/03

October 1995

Do Naked Singularities Form?

CENALO VAZ[†]*Unidade de Ciências Exactas e Humanas**Universidade do Algarve**Campus de Gambelas, P-8000 Faro, Portugal*

and

LOUIS WITTEN^{††}*Department of Physics**University of Cincinnati**Cincinnati, OH 45221-0011, U.S.A.*

ABSTRACT

A naked singularity is formed by the collapse of a Sine- Gordon soliton in 1+1 dimensional dilaton gravity with a negative cosmological constant. We examine the quantum stress tensor resulting from the formation of the singularity. Consistent boundary conditions require that the incoming soliton is accompanied by a flux of incoming radiation across past null infinity, but neglecting the back reaction of the spacetime leads to the absurd conclusion that the total energy entering the system by the time the observer is able to receive information from the singularity is infinite. We conclude that the back reaction must prevent the formation of the naked singularity.

[†] INTERNET: CVAZ@MOZART.SI.UALG.PT

^{††} INTERNET: WITTEN@PHYSUNC.PHY.UC.EDU

Roughly speaking naked singularities are singularities that may be seen by physically allowed observers. So far they have been considered intuitively undesirable and this has produced a conjecture that forbids their existence, the so-called cosmic censorship conjecture. The conjecture simply states that a physically reasonable system of energy will not, under reasonable initial conditions, evolve into a naked singularity. The original intent of the conjecture was of course to prohibit the formation of naked singularities in classical general relativity so that, for example, a non-rotating system would radiate away all multipole moments sufficiently rapidly that the final state would be a Schwarzschild black hole. However, there are many counter examples in the literature where naked singularities do indeed form from classically reasonable initial conditions.¹

The existence of naked singularities is still difficult to understand physically, and several suggestions have been made to maintain the viability of the cosmic censorship hypothesis. One is merely to reiterate that it holds classically but needs some fine tuning to describe the set of physically reasonable set of states from which a collapse may begin. The fine tuning would be such as to exclude all known examples of the formation of naked singularities. A second way to preserve the conjecture is to argue from the point of view of string theory.² String theory is said to solve the short distance problems of general relativity by providing a fundamental length scale proportional to the inverse square root of the string tension. The expectation is therefore that the high energy behavior of string theory (in weak coupling) would forbid any observations of processes associated with high field gradients over short distances. Yet a third possibility, which we have argued for elsewhere,³ is in contradistinction to the string argument. It is that when the singularity forms the accompanying Hawking evaporation is huge and quantum gravity or the back reaction must prevent the singularity from forming.

In this letter we present a model of collapse leading to the formation of a naked singularity and show that, at least in the one loop approximation and without taking the back reaction into account, the Hawking radiation is incoming and infinite. This suggests, we argue, that because of quantum gravity effects or the back reaction the singularity will not form but the energy will dissipate, perhaps explosively. The model we consider is 1+1 dimensional gravity with Sine-Gordon solitons⁴ described by the action

$$S = \frac{1}{2\pi} \int d^2x \sqrt{-g} \left[e^{-2\phi} (-R + 4(\nabla\phi)^2 + \Lambda) - \frac{1}{2}(\nabla f)^2 + 4\mu^2 e^{-2\phi} (\cos f - 1) \right] \quad (1)$$

where ϕ is the dilaton, f is the matter field and Λ is the cosmological constant which we take here to be negative, $\Lambda = -4\lambda^2$. R is the scalar curvature, and $g_{\mu\nu}$ is the two-metric. Our conventions are those of Weinberg.⁵

It is simplest to analyze the field equations in the conformal gauge, where the metric has the form

$$g_{\mu\nu} = e^{2\rho}\eta_{\mu\nu} \quad (2)$$

and in lightcone coordinates, $x^\pm = x^0 \pm x^1$. If we require that the metric returns to the linear dilaton vacuum in the absence of the incoming soliton, the solution to the field equations is given by

$$\begin{aligned} f_{kink} &= 4 \tan^{-1} e^{(\Delta - \Delta_0)} \\ \sigma &= \lambda^2 x^+ x^- - 2 \ln \cosh(\Delta - \Delta_0) \end{aligned} \quad (3)$$

where $\sigma = e^{-2\rho} = e^{-2\phi}$ and $\Delta = \gamma_+ x^+ + \gamma_- x^-$ with

$$\gamma_\pm = \pm \mu \sqrt{\frac{1 \pm v}{1 \mp v}}. \quad (4)$$

Also, v is the velocity of the soliton, $f(x, t) = f(x + vt)$, and $\Delta = \Delta_0$ is its center. We take $\Delta_0 > 0$ throughout, without any loss of generality.

The curvature singularity is at $\sigma = 0$, as can be verified directly by examining the expression ($R = 4\sigma\partial_+\partial_-\ln\sigma$) for the curvature scalar. The Kruskal diagram displayed in figure I shows the singularity along with the trajectory of the soliton center. In this letter we will be particularly interested in the lower branch of the singularity. It is made up of a spacelike piece in the far future joined smoothly to two timelike singularities on either end, being null in the neighborhood of $\mathcal{I}^-$, and intersecting $\mathcal{I}_{\mathcal{R}}^-$ at $x^- = -\infty, x^+ = 2\gamma_-/\lambda^2$ and $\mathcal{I}_{\mathcal{L}}^-$ at $x^+ = -\infty, x^- = -2\gamma_+/\lambda^2$. The upper branch cannot be regarded as a true “formation” of a naked singularity as the soliton energy and singularity appear in the spacetime together, the soliton emerging from the singularity and travelling all the way to i^+ . This case has been examined in ref.[6].

Classical soliton energy entering the spacetime therefore produces a naked singularity in the future. Accompanying the production of the singularity is Hawking radiation. The stress tensor of this radiation is completely determined by the trace anomaly and

the conservation equations. The trace anomaly, in turn, can depend only on the scalar curvature, R , as this is the only curvature invariant with the correct length dimension. The conservation laws determine the stress tensor up to two functions, one of x^+ and the other of x^- according to

$$\begin{aligned}\langle T_{++} \rangle &= T_{++}^f + T_{++}^q = T_{++}^f - \alpha \left[\frac{\partial_+^2 \sigma}{\sigma} - \frac{1}{2} \left[\frac{\partial_+ \sigma}{\sigma} \right]^2 \right] + A(x^+) \\ \langle T_{--} \rangle &= T_{--}^f + T_{--}^q = T_{--}^f - \alpha \left[\frac{\partial_-^2 \sigma}{\sigma} - \frac{1}{2} \left[\frac{\partial_- \sigma}{\sigma} \right]^2 \right] + B(x^-) \\ \langle T_{+-} \rangle &= T_{+-}^f + \alpha \partial_+ \partial_- \ln \sigma,\end{aligned}\tag{5}$$

where α is a positive, dimensionless (spin dependent) constant and $T_{\mu\nu}^f$ is the classical soliton stress energy

$$\begin{aligned}T_{++}^f &= \frac{1}{2}(\partial_+ f)^2 = \frac{4\gamma_+^2}{\cosh^2(\Delta - \Delta_0)} \\ T_{--}^f &= \frac{1}{2}(\partial_- f)^2 = \frac{4\gamma_-^2}{\cosh^2(\Delta - \Delta_0)} \\ T_{+-}^f &= \mu^2(\cos f - 1) = -\frac{2\mu^2}{\cosh^2(\Delta - \Delta_0)}.\end{aligned}\tag{6}$$

Appropriate boundary conditions must now be imposed to determine these arbitrary functions. We expect that the tensor is regular at all points in the spacetime except at the singularity, $\sigma = 0$. Naïvely it may seem that natural conditions are (a) there is no incoming flux of energy other than the soliton's and (b) the Hawking radiation vanishes in the absence of the classical soliton stress energy, i.e., when $\mu = 0 = \Delta_0$. However, one finds that the tensor satisfying these conditions is not regular everywhere. This is because requiring the absence of incoming energy on $\mathcal{I}^-$ (other than the soliton, condition (a)) implies that

$$A(x^+) = -\frac{1}{2x^{+'2}}, \quad B(x^-) = -\frac{1}{2x^{-'2}}\tag{7}$$

where

$$\begin{aligned}x^{+'} &= x^+ - \frac{2\gamma_-}{\lambda^2} \\ x^{-'} &= x^- + \frac{2\gamma_+}{\lambda^2},\end{aligned}\tag{8}$$

As the term in square brackets in (5) is finite everywhere, no incoming flux on $\mathcal{I}^-$ implies that $\langle T_{++} \rangle \rightarrow -\infty$ on the lightlike line $x^{+'} = 0$ and $\langle T_{--} \rangle \rightarrow -\infty$ on the lightlike line

$x^{-'} = 0$. This behavior is physically unacceptable on the grounds that there is nothing special about the spacetime along these lines and infinite negative fluxes to the left and right are in violation of the positive energy conditions.

On the other hand the tensor that is regular everywhere within the spacetime (except at the singularity, $\sigma = 0$) and vanishes in the absence of the soliton is given by

$$\begin{aligned}\langle T_{++} \rangle &= T_{++}^f + T_{++}^q = T_{++}^f - \alpha \left(\frac{\partial_+^2 \sigma}{\sigma} - \frac{1}{2} \left[\frac{\partial_+ \sigma}{\sigma} \right]^2 \right) - \frac{\alpha}{2x^{+2}} \\ \langle T_{--} \rangle &= T_{--}^f + T_{--}^q = T_{--}^f - \alpha \left(\frac{\partial_-^2 \sigma}{\sigma} - \frac{1}{2} \left[\frac{\partial_- \sigma}{\sigma} \right]^2 \right) - \frac{\alpha}{2x^{-2}} \\ \langle T_{+-} \rangle &= T_{+-}^f + \alpha \partial_+ \partial_- \ln \sigma.\end{aligned}\tag{9}$$

In coordinates that are manifestly asymptotically flat

$$\begin{aligned}x^+ &= -\frac{1}{\lambda} e^{-\lambda \sigma^+} + \frac{2\gamma_-}{\lambda^2} \\ x^- &= -\frac{1}{\lambda} e^{-\lambda \sigma^-} - \frac{2\gamma_+}{\lambda^2}\end{aligned}\tag{10}$$

the tensor approaches

$$\begin{aligned}T_{--}^{(\sigma)} &\rightarrow 0, & T_{+-}^{(\sigma)} &\rightarrow 0 \\ T_{++}^{(\sigma)} &\rightarrow \frac{\alpha \lambda^2}{2} \left[1 - \frac{1}{\left(1 - \frac{2\gamma_-}{\lambda} e^{\lambda \sigma^+} \right)} \right]\end{aligned}\tag{11}$$

on $\mathcal{I}_{\mathcal{R}}^-$ and

$$\begin{aligned}T_{++}^{(\sigma)} &\rightarrow 0, & T_{+-}^{(\sigma)} &\rightarrow 0 \\ T_{--}^{(\sigma)} &\rightarrow \frac{\alpha \lambda^2}{2} \left[1 - \frac{1}{\left(1 + \frac{2\gamma_+}{\lambda} e^{\lambda \sigma^-} \right)} \right]\end{aligned}\tag{12}$$

on $\mathcal{I}_{\mathcal{L}}^-$. Therefore consistency seems to require an incoming flux of radiation across past null infinity. This flux is seen to increase smoothly from zero on i^- to the constant value $\alpha \lambda^2/2$ as the lightlike lines $x^- = -2\gamma_+/\lambda^2$ and $x^+ = 2\gamma_-/\lambda^2$ are approached. The total energy flowing into the spacetime is the integrated flux from i^- to the point $x^+ = 2\gamma_-/\lambda^2$ on the right and $x^- = -2\gamma_+/\lambda^2$ on the left. This is obviously infinite, a result that is nonsense. It is easy to see that this will occur whenever the singularity is such as to be

visible to a timelike observer. This is possible in the soliton model because of the timelike part of the singularity in the neighborhood of $\mathcal{I}^-$. As the total energy entering the system across past null infinity cannot be greater than the classical soliton energy, the effect of the incoming Hawking radiation will be to de-localize the soliton. On the other hand, the back reaction can be expected to reduce the spreading of the soliton across $\mathcal{I}^-$. This would be possible only if the back reaction works toward eliminating the timelike piece of the singularity. Indeed we may conclude that quantum- gravity effects, the back reaction of the spacetime, will not permit the formation of the classical singularity as described earlier and that a simultaneous quantization of all fields in the system is necessary for even a qualitative understanding of the process.

The above arguments have been based on what one might physically expect of the back reaction of the spacetime. To incorporate the quantum effects to lowest order in studying the back reaction, one must include the contribution of the conformal anomaly (which arises because the measure on the space of matter fields is non- invariant) in the effective action. An important question is therefore whether the model being considered here is consistent, that is, if the equations of motion that have been solved above are consistent to lowest order. Below we will briefly show, following arguments by de Alwis⁷ and Bilal and Calan⁸, that the Sine-Gordon theory in (1) is actually a special case of such a consistent theory. Our arguments will closely follow those of ref.[7]

We can expect that the full quantum action, including matter but not including ghosts takes the form (in terms of some fiducial metric $g = e^{2\rho}\hat{g}$)⁷

$$S[X, \hat{g}] = - \int \sqrt{\hat{g}} \left[-\frac{1}{2} \hat{g}^{\mu\nu} G_{ab} \nabla_\mu X^a \nabla_\nu X^b - \hat{R}\Phi + T(X) \right] \quad (13)$$

where T is the tachyon potential and X^a is the $N + 2$ dimensional vector

$$X^a = \begin{bmatrix} \phi \\ \rho \\ f_1 \\ \cdot \\ \cdot \\ \cdot \\ f_N \end{bmatrix} \quad (14)$$

including the N matter fields. The modified CGHS theory we have considered is the weak

coupling limit of (13), i.e., in the limit $e^{2\phi} \ll 1$, (13) should look like

$$\begin{aligned}
S[\phi, \rho, f_i] &= \int d^2\sigma \sqrt{\hat{g}} \left[e^{-2\phi} \left(4(\hat{\nabla}\phi)^2 - 4\hat{\nabla}\phi \cdot \hat{\nabla}\rho \right) - \alpha(\hat{\nabla}\rho)^2 \right. \\
&\quad \left. - \hat{R} \left(e^{-2\phi} - \alpha\rho \right) + \Lambda e^{2(\rho-\phi)} \right. \\
&\quad \left. - \frac{1}{2} \sum_i (\hat{\nabla}f_i)^2 + 4\mu^2 e^{2(\rho-\phi)} U(\vec{f}) \right]
\end{aligned} \tag{15}$$

where, in the case we are studying, $U(\vec{f})$ is the Sine-Gordon potential. Comparing (13) and (15), one finds that the low energy limit is given by

$$\begin{aligned}
G_{ab} &= \begin{bmatrix} -8e^{-2\phi} & 4e^{-2\phi} & 0 & . & . & . & 0 \\ 4e^{-2\phi} & 2\alpha & 0 & . & . & . & 0 \\ 0 & 0 & 1 & . & . & . & . \\ 0 & 0 & 0 & 1 & . & . & . \\ . & . & . & . & . & . & . \\ . & . & . & . & . & . & 1 \end{bmatrix} \\
\Phi &= +\alpha\rho - e^{-2\phi} \\
T(X) &= -\Lambda e^{2(\rho-\phi)} - 4\mu^2 e^{2(\rho-\phi)} U(\vec{f}).
\end{aligned} \tag{16}$$

Now consider a generalization of the above field space metric, but with the appropriate limit

$$ds^2 = -8e^{-2\phi}(1+h_1(\phi))d\phi^2 + 8e^{-2\phi}(1+h_2(\phi))d\phi d\rho + 2\alpha(1+h_3(\phi))d\rho^2 + \sum_i df_i^2 \tag{17}$$

where $h_i(\phi)$ are $O(e^{2\phi})$ functionals of the dilaton, ϕ . The metric can be brought to a flat form by the following field redefinitions:

$$\begin{aligned}
y &= \sqrt{2\alpha} \left[\rho - \frac{1}{\alpha} e^{-2\phi} + \frac{2}{\alpha} \int e^{-2\phi} h_2(\phi) d\phi \right] \\
x &= \int P(\phi) d\phi
\end{aligned} \tag{18}$$

where

$$P(\phi) = e^{-2\phi} \left[8e^{2\phi}(1+h_1) + \frac{8}{\alpha}(1+h_2)^2 \right]^{\frac{1}{2}} \tag{19}$$

which give for the metric in (17)

$$ds^2 = -dx^2 + dy^2 + \sum_i df_i^2. \quad (20)$$

In the weak field limit,

$$\begin{aligned} y &\sim \sqrt{2\alpha} \left(\rho - \frac{1}{\alpha} e^{-2\phi} + \dots \right) \\ x &\sim 2\sqrt{\frac{2}{\alpha}} \left(\frac{\alpha}{2} \phi - \frac{1}{2} e^{-2\phi} + \dots \right) \end{aligned} \quad (21)$$

Let us now consider the beta-function equations for the action in (13)

$$\begin{aligned} 0 &= \beta_{ab}^G = -\mathcal{R}_{ab} + 2\nabla_a \nabla_b \Phi - \nabla_a T \nabla_b T + \dots \\ 0 &= \beta^\Phi = -\mathcal{R} + 4G^{ab} \nabla_a \Phi \nabla_b \Phi - 4\nabla^2 \Phi + \frac{N-24}{3} + \dots \\ 0 &= \beta^T = -2\nabla^2 T + 4G^{ab} \nabla_a \Phi \nabla_b T - 4T + \dots \end{aligned} \quad (22)$$

It is easy to see, using (20), that the first of the above equations implies that Φ is a linear function of the fields, and to have the appropriate limit it must be uniquely given by

$$\Phi = \sqrt{\frac{\alpha}{2}} y \quad (23)$$

The second beta function equation gives trivially $\alpha = (N-24)/6$, and the third implies that

$$\partial_x^2 T - \partial_y^2 T + \sqrt{2\alpha} \partial_y T - \sum_i \partial_{f_i}^2 T - 2T = 0 \quad (24)$$

where we have retained only first order terms in the tachyon potential. Consider, first, that $T(X)$ depends only on the fields (x, y) . Obviously, a solution that will reduce to the appropriate one in the limit of weak coupling is

$$T(x, y) = \Gamma e^{\sqrt{\frac{2}{\alpha}}(y-x)} \sim \Gamma e^{2(\rho-\phi)} \quad (25)$$

where Γ is some constant. Including the f fields, if T is of the form

$$T(X) \sim e^{\sqrt{\frac{2}{\alpha}}(y-x)} U(\vec{f})$$

i.e., if $T(x, y, f_i)$ has the same coupling to the ρ and ϕ fields as the cosmological term,

then (24) can be satisfied only if $U(\vec{f})$ is harmonic,

$$\sum_i \partial_{f_i}^2 U(\vec{f}) = 0 \quad (26)$$

If one restricts the number, N , of fields, f_i , to be even, the potential

$$U(\vec{f}) = 4\mu^2 \prod_{i=1}^{N/2} \cos f_i \prod_{j=N/2+1}^N \cosh f_j \quad (27)$$

satisfies the condition (26). Now (13) is linear in T (to this approximation), so

$$T(X) = -\Lambda e^{2(\rho-\phi)} - 4\mu^2 e^{2(\rho-\phi)} \left(\prod_{i=1}^{N/2} \cos f_i \prod_{j=N/2+1}^N \cosh f_j - 1 \right) \quad (28)$$

is a consistent potential. It is also consistent with the equations of motion derived from (1) as taking $f_i = 0$ for $i \geq 2$ gives back the soliton model and, for any f_i , $f_i = 0$ is a solution of the classical equations of motion with this potential.

In conclusion, the following scenario seems to emerge. Ignoring quantum effects, the observer would enter spacetime at i^- accompanied by highly localized soliton energy and doomed to finally crash into a singularity. Before doing so he would have to cross $x^+ = 2\gamma_-/\lambda^2$ or $x^- = -2\gamma_+/\lambda^2$ or both. Once he has crossed these lines he would be able to receive information from the singularity and would realize that the cosmic censorship hypothesis could not hold in his universe. As we have seen, however, quantum effects will play an important role. The stress tensor that is regular throughout the spacetime except at the singularity itself indicates that on entering the spacetime the observer encounters a flux of incoming energy across $\mathcal{I}_{\mathcal{L}}^-$ and $\mathcal{I}_{\mathcal{R}}^-$ and accompanying the soliton. Neglecting the back reaction, the flux appears to increase steadily from i^- leading to an infinite total energy entering the spacetime *before* the singularity becomes actually visible. This is obviously physically inviable and the back reaction can be expected to step in, stop the process and prevent the naked singularity from actually forming.

Acknowledgements

This work was supported in part by NATO under contract number CRG 920096. C. V. acknowledges the partial support of the *Junta Nacional de Investigação Científica e Tecnológica* (JNICT) Portugal under contracts number CERN/S/FAE/128/94 and number CERN/S/FAE/1037/95, and L.W. acknowledges the partial support of the U. S. Department of Energy under contract number DOE-FG02-84ER40153.

References

1. P. S. Joshi, *Global Aspects in Gravitation and Cosmology*, Clarendon Press, Oxford, 1993. Chapter 6.
2. E. Martinec, *Class. Quant. Grav.* **12** (1995) 941
3. C. Vaz and L. Witten, *Phys. Letts.* **B325** (1994) 27.
4. Hak-Soo Shin and Kwang-Sup Soh, *Phys. Rev.* **D52** (1995) 981.
5. S. Weinberg *Gravitation and Cosmology*, John Wiley & Sons, Inc., N.Y., (1972).
6. C. Vaz and L. Witten, *Class. Quant. Grav.* **12** (1995) 2607.
7. S. P. de Alwis, *Phys. Letts.* **B289** (1992) 278; *Phys. Letts.* **B300** (1993) 330; *Phys. rev.* **D46** (1992) 5429.
8. A. Bilal and C. Calan, *Nucl. Phys.* **394** (1993) 73.

Figure Caption:

1. The Kruskal diagram for $\Lambda = -4\lambda^2 < 0$ and $\Delta_0 > 0$. Regions II & IV are physical ($\sigma > 0$). Two timelike singularities are joined smoothly at the soliton center. The figure was drawn for the following parameter values: $\mu = 1$, $\lambda = 0.25$, $\gamma_+ = \sqrt{3}$, $\gamma_- = -1/\sqrt{3}$, $\Delta_0 = 5.0$ and $\alpha = 1/24\pi$.




FIGURE I

